

AD _____

GRANT NUMBER DAMD17-97-1-7044

TITLE: Role of Tissue Transglutaminases in Breast Cancer

PRINCIPAL INVESTIGATOR: Zishan A. Haroon, M.D.

CONTRACTING ORGANIZATION: Duke University
Durham, North Carolina 27710

REPORT DATE: July 1998

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE July 1998	3. REPORT TYPE AND DATES COVERED Annual (1 Jul 97 - 30 Jun 98)		
4. TITLE AND SUBTITLE Role of Tissue Transglutaminases in Breast Cancer		5. FUNDING NUMBERS DAMD17-97-1-7044		
6. AUTHOR(S) Zishan A. Haroon, M.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Duke University Durham, North Carolina 27710		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
11. SUPPLEMENTARY NOTES		19990105 131		
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) Angiogenesis is vital to tumor growth and metastasis. The scope of this grant was to study in detail the role of Tissue transglutaminase (tTG) during wound healing and tumorigenesis. In the first year of the grant proposal, we have described the expression, localization, molecular form and tTG's association with other major determinants of wound healing and tumorigenesis. Our findings clearly show that tTG is readily upregulated in wound healing and rat mammary adenocarcinoma and is associated with endothelial and inflammatory cells. Hypoxia, Vascular endothelial growth factor, Transforming growth factor beta and Tumor necrosis factor alpha are also upregulated alongside tTG in those cells. tTG is quickly proteolysed in the tumor necrosis and that may have important consequences as tTG can hydrolyze ATP/GTP in its fragmented form. We are submitting these findings to Journal of Clinical Investigation, Proceedings of National Academy of Sciences and American Journal of Pathology where we have detailed the potential significance of these observations. We have also addressed most of the tasks detailed in the statement of work and are now set to proceed with the rest of the project.				
14. SUBJECT TERMS Breast Cancer, Angiogenesis, Tissue Transglutaminase, Fibrin, Extracellular Matrix		15. NUMBER OF PAGES 8		
		16. PRICE CODE		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

X In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

X In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

 7/30/98

PI - Signature Date

TABLE OF CONTENTS

Cover

SF 298 (Report Documentation Page)

Foreword

Table of Contents

Introduction 5

Body 5

Conclusions 6

References 7

List of Publications 8

INTRODUCTION:

Tissue Transglutaminase (tTG) is a member of family of important enzymes called Transglutaminases. These are calcium dependent enzymes which catalyze intermolecular covalent bonds. This crosslinking makes the proteins resistant to mechanical and proteolytic degradation. The purpose of our grant proposal was to describe the role of this extracellular matrix enzyme during wound healing and tumorigenesis as both processes are very similar in mechanism.

The importance of tTG in wound healing was initially based on the presence of defective wound healing in factor XIII deficiency. Bowness et al described the presence of tTG in wound healing in rats and made the important observation that although antigen and activity both increased, there was more antigen than activity indicating degradation or inactivation of the protein [1]. Application of putrescine, a transglutaminase inhibitor, produced a significant decrease in wound strength between day 5 and 10 [2].

tTG's involvement in tumor biology has been demonstrated by numerous studies. An inverse relationship between tTG activity and metastatic potential has been reported by several investigators in cells of murine origin [3,4]. Knight et al reported on direct relationship between tTG activity and detergent insoluble apoptotic body formation in a number of metastatic cell lines cloned from a hamster fibrosarcoma [5]. Johnson et al transfected human tTG cDNA hamster sarcoma Met B cells and reported that overexpression of tTG lead to delay and suppression of tumor cell growth [6].

In addition, tTG has been implicated in apoptosis, cellular signaling, bone formation and its list of substrates is extensive and includes all major extracellular proteins. Thus, it is vital to understand the basic mechanisms regarding the working of this enzyme to effectively use it for novel therapeutics for breast cancer.

BODY:

We realized the need of first describing the expression, molecular form of tTG and its association with other major modulators of wound healing such as Vascular Endothelial Growth Factor (VEGF), Transforming Growth Factor beta (TGF beta), Tumor Necrosis Factor alpha (TNF alpha) and Hypoxia. This would clearly help establish tTG's role during wound healing and tumorigenesis and also suggest possible mechanisms to its role during these processes.

We are now in the process of submitting this information as three publications (see appendix) which detail our findings in this regard. We have found that tTG is expressed by endothelial cells, macrophages and fibroblasts

during both wound healing and R3230 Mammary adenocarcinoma. tTG expression is most closely related to TGF beta. tTG is also rapidly degraded and is present in 50-55 kda and 20 kda form. We will forward the final accepted versions of the papers later.

We have also addressed the major tasks to be completed during the first year. We will detail them task by task:

Task 1 and 3: We have isolated fibrinogen from rats and prepared for the experiments outlined in the grant. We have also ran comparisons of that fibrin with the commercially available fibrin from Sigma and found that our isolation procedure deactivates native Factor XIIIa, which results in weak fibrin gels. We will supplement the gels with 5ug/ml of Factor XIIIa for the experiments.

We also have expressed recombinant tTG and its mutants in a E. Coli based expression system. Our yield of the protein was low in earlier expressions and we had to repeat the procedure to have enough quantity of the proteins for future experiments.

Both these processes took a little more time than expected which pushed our time line on other tasks.

Task 5 and 6: We started by manufacturing our own chambers but we could not reduce the size of the chamber down from 3 mm in width. We have now moved on with Millipore chambers and modified them for our use. The width is down to 1.5 mm which will save a lot of precious proteins and time from now on.

We are in the process of doing fibrin gel experiments to assess the dose curve of tTG in the fibrin chamber. We will relate the data in next report.

Task 2: We are setting up fibrinolysis assays to assess the tTG's ability to impart more ability to the fibrin gel and make it more resistant to proteolysis against plasmin. In this regard, I am also collaborating with other investigators in using a rat aorta model developed by Nicosia to better answer this question.

Task 4: We have not been able to address this aim in the last year due to overruns by other tasks. we have started work on this aim and are In the process of establishing the required assays to accomplish this task.

CONCLUSIONS:

Our work has established tTG as one of the prominent players during wound healing and tumor development and also how it is associated with other prominent cytokines and cells during these processes. Its ability to enhance angiogenesis during wound healing and limit tumor growth gave interesting insights into its activities. This sets up nicely for our work with fibrin gel experiments which would define the mechanisms of its effects during wound healing and tumor development.

REFERENCES:

1. J M Bowness,, A H Tarr, T Wong ¶: Increased Transglutaminase Activity During Skin Wound Healing in Rats. *Biochimica et Biophysica Acta*. 967: 234-240 (1988).
2. K N Dolynchuk, R Bendor-Samuel, J M Browness : Effect of Putrescine on Tissur Transglutaminase Activity in Wounds: Decreased Breaking Strength and Increased Matrix Fucoprotein Solubility. *Plastiic and Reconstructive Surgery*. 93: 567-573 (1994).
3. Beninati, S., Abbruzzese, A., and Cardinali, M., Differences in the post-translational modification of proteins by polyamines between weakly and highly metastatic B16 melanoma cells. *Int J Cancer*,. 53(5): p. 792-7, 1993.
4. Hand, D., Elliott, B.M., and Griffin, M., Expression of the cytosolic and particulate forms of transglutaminase during chemically induced rat liver carcinogenesis. *Biochim Biophys Acta*, 970(2): p. 137-45, 1988.
5. Knight, C.R., Rees, R.C., and Griffin, M., Apoptosis: a potential role for cytosolic transglutaminase and its importance in tumour progression. *Biochim Biophys Acta*, 1096(4): p. 312-8, 1991.
6. Johnson, T.S., Knight, C.R., el-Alaoui, S., Mian, S., Rees, R.C., Gentile, V., Davies, P.J., and Griffin, M., Transfection of tissue transglutaminase into a highly malignant hamster fibrosarcoma leads to a reduced incidence of primary tumour growth. *Oncogene*, 9(10): p. 2935-42, 1994.

List Of Publications/Abstracts where Defense Grant was Cited:

1. **ZA Haroon**, KG Peters, CS Greenberg and MW Dewhirst. Ch:1 Angiogenesis and oxygen transport in the solid tumors in *Antiangiogenic Agents* (BA Teicher, ed) Humana, Totowa, NJ (In press).
2. **ZA Haroon**, JM Hettasch, TS Lai, RL McCauly, MW Dewhirst and CS Greenberg. Tissue Transglutaminase expression during rat dermal wound healing and it can function as a pro-angiogenic molecule (Being submitted to Journal Of Clinical Investigation)
3. **ZA Haroon**, JA Raleigh, CS Greenberg and MW Dewhirst. Interrelationship between Hypoxia, Cytokines and Inflammatory cells during rat dermal wound healing. (Being submitted to Proceedings of National Academy of Sciences)
4. **ZA Haroon**, R Syal, RL McCauly, RA Lindberg, JM Hettasch, MW Dewhirst and CS Greenberg. Tissue Transglutaminase blocks growth of R3230 Mammary Adenocarcinoma in Rat window chamber (Being submitted to American Journal Of Pathology).
5. MR Matthews, **ZA Haroon**, AH Friedman, CJ Wikstrand, CS Greenberg, JL Burchette, DD Bigner, RE McLendon. Extracellular Matrix Antigens in Capillary Hemangioblastoma: an Immunohistochemical Study. (Being submitted to Journal of Histochemistry and Cytochemistry).

ZA Haroon, JM Hettasch, MW Dewhirst, CS Greenberg. The Transglutaminase Enzyme Derived from Vascular Cells Contributes To the Proangiogenic Properties of Fibrin. *39th Annual ASH, California, December, 1997.*

ZA Haroon^{e*}, RA Lindberg, R Syal, CS Greenberg, MW Dewhirst. Localization of Macrophages in Tumor has Implications Regarding Tumor Angiogenesis Stroma Formation. *AACR Special Conference in Cancer Research, Florida, January, 1998.*

KL Blackwell, **ZA Haroon**, JK Parker, MW Dewhirst, and CS Greenburg. Serum VEGF correlates wit breast tumor size but not with patterns of immunohistochemically detected VEGF deposition. *AACR Special Conference in Cancer Research, Florida, January, 1998.*

RL Lindberg, **ZA Haroon**, TS Lai, JM Hettasch, MW Dewhirst and CS Greenburg. Inhibition of PMN adhesion to endothelial cells by tissue transglutaminase. *AACR Annual Meeting, Loiusiana, March, 1998.*

Salary Support:

Zishan A. Haroon MD